Space Weather Technical Forecast

Issued on Monday, 14 August 2023 at 13:14 Local

This technical forecast provides a four day assessment of space weather events. The probabilities stated below are for reaching or exceeding the given levels. For more information about space weather impacts please see the Met Office Space Weather Scales https://www.metoffice.gov.uk/weather/learn-about/space-weather/uk-scales

Space Weather Forecast Headline: Chance of G1/Minor Storm, mainly day 2 (15 Aug). Chance of isolated M-flares (R1 Blackouts).

Analysis of Space Weather Activity over past 24 hours

Solar Activity: Low with peak flare C2.0 from AR3395 at 13/2232 UTC. There are four numbered, and three unnumbered regions on the visible disk. AR3403 in the northeast remains the most developmental, however this has now settled into a relatively simple bipolar configuration. AR3395 in the northwest, has continued to slowly weaken but remains magnetically complex for now. AR3397 is a simple open bipolar spot. AR3394 is a mature and stable unipolar spot, now approaching the southwest limb. AR3402 faded to plage. The other regions are smaller and simple. Further regions appear about to rotate onto the east limb. There were no Earth-directed CMEs.
Solar Wind / Geomagnetic Activity: Slow wind conditions observed by ACE at L1 with wind speeds around 350 km/s. Density remained below average. Total interplanetary magnetic field (IMF) Bt, was weak, below 6nT, with north-south component, Bz, varying between -4 nT and +4nT. Phi angle was predominately negative (towards the Sun) before rotating into a positive orientation around 14/0900UTC. Geomagnetic activity was Quiet (Kp 1-2).
Energetic Particles / Solar Radiation: GOES16 high energy (greater than 10MeV) proton flux has remained at background levels. GOES16 High energy (greater than 2MeV) electron flux varied between background and high levels with the diurnal cycle, peaking 1370 pfu at 13/1530 UTC. A slower decline was observed than recent days, however. The associated 24-hour integrated fluence remained below Active (1e8 integrated pfu), with a steady trend, reaching 5.91e7 integrated pfu at 14/0000UTC.

Four-Day Space Weather Forecast Summary
Solar Activity: Mainly Low activity is expected, but with a daily chance of rising to Moderate with isolated M-flares.
Solar Wind / Geomagnetic Activity: There are no Earth-directed CMEs. CH33/+ fast winds are anticipated to arrive either later day 1 or more likely day 2 (14-15 Aug), with solar winds likely rising to around 500-600km/s. Further enhancements are possible day 3 (16 Aug) from CH35/-, however this is low confidence. Initially Quiet geomagnetic activity is expected to become Unsettled to Active with a likelihood of G1/Minor Storm intervals, either late day 1 or day 2 (14-15 Aug). Activity then easing to Unsettled to Active, but with an ongoing slight chance of G1 Minor Storm intervals day 3 and 4 (16-17 Aug).
Energetic Particles / Solar Radiation: High energy (greater than 10 MeV) proton flux is expected to persist at background. High energy electron flux (greater than 2MeV) is forecast to continue varying between Background and High, but with an increasing trend over previous days, before dropping out in response to any fast wind arrival later day 1 or day 2 (14-15 Aug). A more persistent recovery to Moderate to High values is then expected follow, however the timing of this flux being observed at GEO is low confidence. Fluence will rise in response, giving a chance of reaching Active, mainly day 3-4 (16-17 Aug).

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Figure 1. Solar Analysis Valid 14/0700 UTC.



.

Key: Filament ___ , Prominence ___ , Magnetic Field Line - - - , Polarity +/-, Coronal Holes: Grey shaded area CHxx +/-, Sunspot groups 25xx - Mt Wilson α-β-βγ-βγδ and Zurich-McIntosh Axx etc.

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Geomagnetic Storms:

There are no Earth-directed CMEs.
Current slow wind conditions are forecast to increase later day 1 (14 Aug) and more likely during day 2 (15 Aug). This is due to the onset of fast winds from CH33/+ in the northern disc, along with any preceding CIR. As a positive polarity coronal hole, there is a higher chance of a geoeffective connection at this time of year (Russel-McPherron effect). This is also a later onset than MOSWOC Enlil indicates, as the model appears to be also generating a high speed stream from an area of weak fields currently in the west centre disc which is currently considered as only having a low chance of occurring earlier on day 1 (14 Aug).
CH35/- on the southern disc has a less certain influence on the expected solar wind conditions, due its negative polarity, and with Earth currently above the ecliptic. This may prolong elevated solar winds into day 3 (16 Aug), but with a lower likelihood of significantly enhanced geomagnetic activity. A gradual easing of winds is then expected day 4 (17 Aug).
Initially Quiet activity is expected to become Unsettled to Active with a likelihood of G1/Minor Storm intervals, either late day 1 or day 2 (14-15 Aug). Activity then easing to Unsettled to Active, but with an ongoing slight chance of G1 Minor Storm intervals day 3 and 4 (16-17 Aug).

Geo-Magnetic Storm Probability (Exceedance)	Level	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC) (%)	Day 2 (00-24 UTC) (%)	Day 3 (00-24 UTC) (%)	Day 4 (00-24 UTC) (%)
Minor or Moderate	G1 to G2	No	30	60	20	10
Strong	G3	No	1	1	1	1
Severe	G4	No	1	1	1	1
Extreme	G5	No	1	1	1	1

Geomagnetic Activity - Earthbound Coronal Mass Ejections

Figure 2: Estimated Planetary K index (Kp) from BGS.



Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Met Office Space Weather Technical Forecast

Radio Blackouts - X-Ray Flares:

Solar activity is currently Low with four numbered sunspot regions and three further small unnumbered regions. There remains a chance of isolated M-flares.
The most active region on the disc is AR3395. This is a magnetically complex, but small and weakening region with only C-flares observed. AR3403 in the northeast is now the main flare risk on the disc, having rapidly developed in the last 24 hours. However this region remains relatively simple with a straightforward bipolar magnetic configuration. The other numbered regions are AR3397 in the northwest which has also decreased in size recently, and AR3394, currently the largest spot on the disc which is now approaching the southwest limb as a mature and stable region. The other regions on the disc are small, simple and unlikely to produce notable flares.

X Ray Flares	Level	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability			(%)	(%)	(%)	(%)
Active	R1-R2 M Class	No	30	40	40	40
Very Active	R3 to R5 X Class	No	1	1	1	1

Solar Radiation Storms - (High Energy Protons):

The flux of high energy (greater than 10 MeV) is at background. The sunspots on the visible disc, including those on the western hemisphere, are currently relatively simple and currently considered unlikely to produce proton events. Barring any further development of these, the proton flux is expected to remain at background.

Radiation Storms	Level (cm^{-2} sr^{-1} s^{-1})	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability (Exceedance)			(%)	(%)	(%)	(%)
Active	≥ S1	No	2	2	2	2
Very Active	≥ S3 *	No	1	1	1	1
* S3 ≥ 10 MeV ≥ 1000 pfu and / or ≥ 50 MeV ≥ 10 pfu. (pfu = $cm^{-2}sr^{-1}s^{-1}$)						

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

High Energy Electrons Event (≥ 2MeV):
The high energy (greater than 2 MeV) electron flux at GOES-16 has been varying between background and diurnal peaks of high (above 1000 pfu). The resulting associated 24 hour fluence has now stabilised from earlier rises and is now on a flat trend. The arrival of the next fast wind on the day 1, or more likely day 2 (14-15 Aug) will lead to the flux dropping out, before a more persistent rise is expected day 3 (onward).
The associated fluence may approach the Active (1e8 integrated) threshold later on day 1 (14 Aug), but most likely staying below, before falling day 2 (15 Aug) as a result of the flux dropping out with any HSS arrival. Day 3 and 4 will then see fluence rising again, with an increasing risk of reaching Active. This is low confidence however, with further solar wind enhancement possible day 3 (16 Aug). REFM is currently reasonable guidance with a steady trend indicated, however a rising or steady trend is considered most likely.

GEO Electron Environment	Level (cm^{-2} sr^{-1} day^{-1})	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability (Exceedance)			(%)	(%)	(%)	(%)
Active	≥ 2 MeV ≥ $1x10^8$	No	10	10	30	40
Very Active	≥ 2 MeV ≥ $1x10^9$	No	1	1	1	1

Figure 3: GOES16 ElectronFlux- Reaching High (1000pfu) at the diurnal maximum each day.



Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Figure 4: MOSWOC REF Model



Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.